

# Intravenous Laser Blood Irradiation in Sports Medicine

Dr. med. F. RAGGI and Dr. med. G. VALLESI, Terni, Italy

Die intravenöse Laserblutbestrahlung ist seit mehr als 20 Jahren bekannt und wurde ursprünglich in Russland entwickelt. Sie wurde vornehmlich zur Behandlung rheumatischer, kardiovaskulärer, pulmonaler und neurologischer Erkrankungen eingesetzt.

Bei dieser Methode wird Laserenergie direkt in das Blut geführt. So entstand die Idee zu überprüfen, ob die intravenös applizierte Lichtenergie die Leistungsfähigkeit von Athleten in der Sportmedizin zu steigern vermag. In dieser kleinen Studie konnte gezeigt werden, dass die intravenöse Laserblutbestrahlung mit

rotem und grünem Laserlicht zu einem lang anhaltenden Effekt auf Kraft und Ausdauer bei verschiedenen Sportarten führt. Eine Hypothese für diesen Effekt könnte eine Aktivierung der Erythrozyten mit erhöhter Sauerstoffbeladung darstellen. Weiterhin wird vermutet, dass der lang anhaltende Effekt über 3-4 Monate mit der mittleren Überlebenszeit der Erythrozyten korreliert sein könnte.

Schlüsselwörter: Intravenöse Laserblutbestrahlung, roter und grüner Laser, Sportmedizin, Erythrozytenaktivierung und -überlebenszeit

## Introduction

Our idea to test the application of intravenous laser blood irradiation in sports medicine started in July 2007, when a young patient of ours told us he was having surprising effects since he had started the treatment.

He was a male, aged 34; he had come to our medical center for a problem of tinnitus. Illness was started 1 year before and he had undergone many diagnostic examinations (MRI, doppler, CAT), but they were all negative. In the previous year, he had had a three months treatment with cortisone, without any success. He was not taking any drugs nor any kind of therapy at the moment we visited him.

The following treatment protocol was used:

- Intravenous Laser
- Laser shower (locally)
- Lipoic Acid (400 mg/die) and Ascorbic acid (100 mg/die), in order to enhance cellular metabolism.

After a 10 sessions treatment, tinnitus was lowered (about 50%), but did not disappear. Nevertheless, the patient experienced some interesting and unexpected




Fig. 1–3: Pectoral Muscle maximum lifting power, Weight lifting from the floor, Squat

“side effects”: he told us he was feeling less tired in the evening, that his sleeping was improved and he referred a better concentration during the day. But, what is very important, is that he was an athlete (body building) and he had noticed that his maximum lifting power was increased, the running time in the training session was doubled and the number of swimming lanes in the training session was increased. Also his trainer was surprised by his improvements.

Many scientific papers have been written so far, showing biological actions and therapeutic properties of intravenous laser blood irradiation; this new medical technology is currently used for the treatment of different pathologies like rheumatic, cardiovascular, pulmonary and neurological diseases. But, to our knowledge, no scientific works have been performed, so far, about laser blood irradiation in sports medicine. So, we decided to perform a short study, in order to confirm the effects that had been observed in our patient.

## Materials and methods

### Study population

4 male body building athletes were enrolled in the study. Mean age was 23,2.

### Treatment protocol

All athletes underwent 10 sessions of intravenous laser blood irradiation, with the following treatment schedule:

- 630 nm, 2 mW, 20 min. (first session);
- 630 nm, 2 mW, 20 min. + 532 nm, 2 mW 10 min. (9 sessions) .

3 sessions per week were performed. All athletes were invited to avoid any drugs or vitamins during the period of study.

**Outcomes**

The following outcomes were measured:

– Maximum strength tests:

1. Pectoral muscle maximum lifting power. Weight is lifted in the supine position, while athlete lies on the bench (Fig. 1). Maximum weight had to be confirmed by 8 following lifting of sub-maximal weight (80% of maximum).
2. Weight lifting from the floor. Weight is on the floor and athlete has to lift it from standing position (Fig. 2). Maximum weight had to be confirmed by 8 following lifting of sub-maximal weight (80% of maximum).
3. Squat. Weight is put on athlete’s shoulders and athlete has move down and up, once. Maximum weight had to be confirmed by 8 following lifting of sub-maximal weight (80% of maximum).

**Endurance tests:**

1. Nr. of swimming lanes. Is the maximum number of swimming pool lanes that athlete could perform until he felt tired. This test is not time dependent.
2. Cord jumping. Is the maximum time that athlete could perform in cord jumping until he felt tired

**Measurements:**

All athletes were evaluated in 8 different times of the period of study:

1. before the treatment,
2. after the treatment,
3. every 4 weeks after the end of the treatment.

**Results**

Mean pectoral muscle maximum lifting power at any time of measurement is shown in Fig. 4. Mean value seems to increase immediately after the treatment and




Fig. 4: Mean pectoral muscle maximum lifting power (kg) at any time of measurement

this effect seems to last for 16 weeks; afterwards, it seems to disappear and previous condition is restored. This trend is shown in Fig. 5, where mean pectoral muscle maximum lifting power percentage variation is reported. Very similar results were obtained in all the other tests that were performed (See Figg. 6–9).




Fig. 5: Mean pectoral muscle maximum lifting power percentage variation at any time of measurement




Fig. 6: Weight lifting from the floor: mean percentage variation at any time of measurement




Fig. 7: Squat: mean percentage variation at any time of measurement




Fig. 8: Number of swimming pool lanes: mean percentage variation at any time of measurement




Fig. 9: Cord Jumping time: mean percentage variation at any time of measurement

## Discussion

In our study, intravenous laser treatment seemed to be effective in sports medicine, even though we have to highlight that the number of cases was very small. Furthermore, there is big lack of data from the literature on this topic and further studies are needed in this field.

In all the athletes in the study, the muscular empowerment appeared to last for 16 weeks after the end of the treatment. Despite this datum have to be confirmed in bigger studies, we could hypothesize that it is related to the erythrocytes' mean living time: 12–15 weeks. We can imagine that, after a complete cycle of treatment, all circulating red blood cells are "activated"; since they have different "ages", they will survive for different times; but, in any case, the youngest among them will disappear from circulating blood after about 15 weeks and, after that, the therapeutic effect will be lost.

In order to verify the plausibility of our findings, we wanted to perform a deep review of the medical literature on Low Level Laser Therapy (LLLT). The review was conducted looking for "low level laser blood irradiation" in the Pub Med Library. Only articles related to the topic of sports medicine were considered.

First of all, we could mention a polish study (1), showing higher exercise capacity, longer exercise time and longer distance of 6 minutes walk test, in 39 coronary artery disease patients, treated with LLLT on the chest area.

In a second study (2), isolated rat tibial anterior muscle was fatigued with electric current until initial strength was lowered to 50%. Then, the rats were divided into two groups: LLLT treated group and a control group. Laser irradiated group had significantly longer time to reach 50% reduction of strength, significantly higher peak force and significantly lower creatine kinase blood levels (muscle damage index). So, we can state that the outcomes observed in our study could be considered plausible, if compared to other studies showing biological properties of LLLT.

But the way these effects are achieved still remains unclear.

Many studies show various biological effects of laser irradiation, related to our topic.

A very important effect is the blood flow increasing. A Japanese research group (3), performed LLLT on common carotid artery area: blood flow volume in central retinal artery and ophthalmic artery increased.

A similar experiment was carried out by Makihara et al (4): LLLT on right temporomandibular joint area

yield expansion of blood vessels and increase in blood flow volume in superficial temporal artery; surprisingly, this effect was observed bilaterally.

Wasik et al (5), observed PO<sub>2</sub> and SaO<sub>2</sub> increasing, after whole blood sample irradiation with 632 nm laser.

Other important referred effects are erythrocytes' deformability increasing (6–8) and microcirculation increasing (9–11).

But which are the possible molecular mechanisms, explaining the above cited effects?

Mittermayr et al (12) showed that blue laser releases Nitric Oxide (NO) from Haemoglobin (Hb). NO-Hb is a form of met-Hb, with low O<sub>2</sub> linking power. So, free NO released from Hb is a powerful molecule producing vasodilatation and perfusion enhancement; at the same time, Hb O<sub>2</sub> linking power is increased.

Besides, irradiation of monocytes leads to increased NO levels from a preformed store, without activation of NO synthase and releasing of endothelial NO (13).

In a study performed by Simonian et al (14), isolated erythrocyte membrane was irradiated and Cit b 558 III activity was measured; Cit b 558 III is a membrane enzyme that restores met-Hb (H<sup>+</sup>-Hb, CO<sub>2</sub>-Hb, DPG-Hb) to normal Hb, with higher O<sub>2</sub> linking power. Cit b 558 III activity was enhanced after laser irradiation.

In conclusion, the data from our literature review seem to confirm the plausibility of our results, even though obtained from a very small study population. We hope that our preliminary data on intravenous laser therapy and sports medicine will contribute to

Intravenous laser blood irradiation is well known for more than 20 years and was originally developed in Russia. It was used for a lot of different diseases like rheumatic, cardiovascular, pulmonary and neurological indications. This method provides laser energy directly in the blood. So the idea was to prove if the intravenous light energy will increase the performance of athletes in sports medicine. In this small study it could be shown that intravenous laser blood irradiation with red and green laser light leads to a long lasting effect on muscular strength and endurance in different sport activities. It is hypothesized that erythrocytes will be activated with a higher capacity of oxygen transport. Furthermore it is assumed that the long lasting effect of 3–4 months is correlated with the mean survival time of the red blood cells.

**Key words:** Intravenous laser blood irradiation, red and green laser, sports medicine, erythrocyte activation and survival time

focus the attention of many researchers on this interesting topic, so that deeper scientific knowledge could be achieved. Especially for the last question that still is without an answer: why the effects of blood irradiation are systemic and not only local?

## References

- [1] Zyciński P, Krzemińska-Pakuła M, Peszyński-Drews C, Kierus A, Trzos E, Rechciński T, Figiel L, Kurpesa M, Plewka M, Chrzanowski L, Drozd J. *Laser biostimulation in end-stage multivessel coronary artery disease-a preliminary observational study*. *Kardiol Pol*. 2007 Jan;65(1):13–21.
- [2] Lopes-Martins RA, Marcos RL, Leonardo PS, Prianti AC Jr, Muscará MN, Aimbire F, Frigo L, Iversen VV, Bjordal JM. *Effect of low-level laser (Ga-Al-As 655 nm) on skeletal muscle fatigue induced by electrical stimulation in rats*. *J Appl Physiol*. 2006 Jul;101(1):283–8. Epub 2006 Apr 20.
- [3] Mii S, Kim C, Matsui H, Oharazawa H, Shiwa T, Takahashi H, Sakamoto A. *Increases in central retinal artery blood flow in humans following carotid artery and stellate ganglion irradiation with 0.6 to 1.6 microm irradiation*. *J Nippon Med Sch*. 2007 Feb;74(1):23–9.
- [4] Makihara E, Masumi S. *Blood Flow Changes of a Superficial Temporal Artery Before and After Low-level Laser Irradiation Applied to the Temporomandibular Joint Area*. *Nihon Hotetsu Shika Gakkai Zasshi*. 2008 Apr;52(2):167–70.
- [5] Wasik M, Gorska E, Modzelewska M, Nowicki K, Jakubczak B, Demkow U. *The influence of low-power helium-neon laser irradiation on function of selected peripheral blood cells*. *J Physiol Pharmacol*. 2007 Nov;58 Suppl 5(Pt 2):729–37.
- [6] Xian-Qiang Mi, Ji-Yao Chen, Zi-Jun Liang, Lu-Wei Zhou. *In Vitro Effects of Helium-Neon Laser Irradiation on Human Blood: Blood Viscosity and Deformability of Erythrocytes*. *Photomedicine and Laser Surgery*
- [7] Iijima K, Shimiyama N, Shimiyama M, Mizuguchi T. *Effects of low He-Ne laser on deformability of stored human erythrocytes*. *J Clin Laser Med Surg*. 1993 Aug;11(4):185–9.
- [8] Zaleskaya, G.; Sambor, E. *Interaction of Low-Intensity Laser Radiation with Blood and Its Components*. *Journal of Applied Spectroscopy*, Volume 72, Number 2, March 2005, pp. 242–248(7)
- [9] Burduli NM, Piliieva NG. *The dynamics of microcirculation parameters in patients with pneumonia receiving intravenous irradiation of blood as a part of complex treatment*. *Klin Med (Mosk)*. 2007;85(7):48–50.
- [10] Ihsan FR. *Low-level laser therapy accelerates collateral circulation and enhances microcirculation*. *Photomed Laser Surg*. 2005 Jun;23(3):289–94.
- [11] Maegawa Y, Itoh T, Hosokawa T, Yaegashi K, Nishi M. *Effects of near-infrared low-level laser irradiation on microcirculation*. *Lasers Surg Med*. 2000;27(5):427–37.
- [12] Mittermayr R, Osipov A, Piskernik C, Haindl S, Dungal P, Weber C, Vladimirov YA, Redl H, Kozlov AV. *Blue laser light increases perfusion of a skin flap via release of nitric oxide from hemoglobin*. *Mol Med*. 2007 Jan–Feb;13(1–2):22–9.
- [13] Lindgård A, Hultén LM, Svensson L, Soussi B. *Irradiation at 634 nm releases nitric oxide from human monocytes*. *Lasers Med Sci*. 2007 Mar;22(1):30–6. Epub 2006 Nov 21.
- [14] Simonian RM, Simonian GM, Simonian MA, Sekoian ES. *Effect of low intensive helium-neon laser on superoxide-producing and methemoglobin-restoring activity of cytochrome b558III of red cell membranes in experiment*. *Vopr Kurortol Fizioter Lech Fiz Kult*. 2007 Jul–Aug;(4):9–11.

Anschriften der Verfasser

Dr. Francesco Raggi, Dr. Giuseppe Vallesi,

Strada di Santa Filomena 16,

05100 Terni, Italien

Tel.: 0039/3285743276,

e-mail: raggimail@virgilio.it, gipovallesi@hotmail.com